NAVAL HEALTH RESEARCH CENTER SAN DIEGO CALIF
SUBJECTIVE EVALUATION OF THE EFFECTS OF SLEEP LOSS: THE NPRU MO--ETC(U)
1974 J M MOSES, A LUBIN, P NAITOH
74-25 AD-A038 367 UNCLASSIFIED OF | AD A038367 END DATE FILMED


( NW


SUBJECTIVE EVALUATION OF THE EFFECTS OF SLEEP LOSS:

THE NPRU MOOD SCALE

- J. MOSES
- A. LUBIN
- P. NAITOH
- L. C. JOHNSON

REPORT NO. 74-25






NAVAL HEALTH RESEARCH CENTER


SAN DIEGO, CALIFORNIA 92152

NAVAL MEDICAL RESEARCH AND DEVELOPMENT COMMAND
BETHESDA, MARYLAND

DISTRIBUTION STATEMENT A

Approved for public release; Distribution Unlimited





SUBJECTIVE EVALUATION OF THE EFFECTS OF SLEEP LOSS: THE NPRU MOOD SCALE

J. M. Moses, A. Lubin, P. Naitoh, and L. C. Johnson

This research was supported in part by National Science Foundation Grants GB-6008 and GB-14829 to the San Diego State University Foundation, and by Department of the Navy, Bureau of Medicine and Surgery, under Task No. MF12.524.004-9008DASG. The opinions and assertions contained herein are the private ones of the writers and are not to be construed as official or as reflecting the views of the Navy Department.

Address requests for reprints and test forms to: Julie M. Moses,
Navy Medical Neuropsychiatric Research Unit, San Diego, California 92152.



DISTRIBUTION STATEMENT A
Approved for public release;
Distribution Unlimited

## Abstract

A mood scale consisting of 52 adjectives and phrases was given to twelve young naval recruits to determine which items were sensitive to at least one night of sleep loss. The items which were maximally sensitive to sleep loss were then used to measure the effects of one night of total sleep loss in another group of fourteen young naval recruits. The two experiments resulted in a list of 29 adjectives and short phrases, the NPRU Mood Scale. This scale can be easily, briefly, and repeatedly employed to detect cumulative sleep loss and to assist in evaluating task performance decrement.

DESCRIPTORS: Adjective check list, Mood scale, Sleep loss,
Task performance. (J. M. Moses)

# SUBJECTIVE EVALUATION OF THE EFFECTS OF SLEEP LOSS: THE NPRU MOOD SCALE

J. M. Moses, A. Lubin, P. Naitoh, and L. C. Johnson

Most sleep deprivation studies have been concerned primarily with the evaluation of performance and of the quality and quantity of sleep, but the development of sensitive valid psychological tests of the subjective effects of sleep loss has been given less attention.

An early study (Murray, Williams, & Lubin, 1958), in which Ss rated themselves on a four-point scale, found significant increases in self-ratings of sleepiness and fatigue, but only after two or three nights without sleep. Recently, Hoddes, Dement, and Zarcone (1972) published the Stanford Sleepiness Scale (SSS). The SSS, which consists of seven statements representing decreasing levels of arousal, is sensitive to one night of sleep loss (Hoddes, Zarcone, Smythe, Phillips, & Dement, (1973).

Thayer (1967, 1970, 1971) and others (e.g., Hendrick & Lilly, 1970; Bohlin & Kjellberg, 1973) have developed and factor analyzed Multiple Adjective Check Lists (MACLs) designed to measure changes in activation, stress, contentment, depression and hostility. These studies reported significant changes in at least one factor following one night of sleep loss.

17 11846

The purpose of this paper is to describe a self-rating MACL, the NPRU Mood Scale. It is a brief and easily administered list of 29 adjectives and short phrases designed to measure mood changes due to one or more nights of sleep loss.

The need for this short test became apparent to us after analysis of the mood data in our recent study of sleep stage deprivation and total sleep deprivation (Johnson, Naitoh, Moses, & Lubin, 1974). In that study, several self-report mood questionnaires were used: an early 52-item version of the Profile of Mood State (POMS) by McNair, Lorr, and Droppleman (1971); the current 65-item version of the POMS; the Primary Affect Scale (PAS) (Johnson & Myers, 1967); and the Spielberger State-Trait Anxiety Inventory (STAI) (Spielberger, 1968). Despite the considerable overlap among these tests, the usual scoring methods produced inconsistent and sometimes contradictory results in describing mood changes after sleep loss.

### METHOD

The NPRU Mood Scale was developed from an experiment (Lubin, Moses, Johnson, & Naitoh, 1974) in which twelve naval recruits, ages 17-21, were deprived of two nights of sleep after a baseline of four days. The original 52-item adjective check list (the early version of the POMS) from which the NPRU Mood Scale was derived was administered at the same time twice each day. The subjects chose one of four possible responses to each item ranging from "not at all" (scored as 0) to "extremely" (scored as 3). Scores were obtained on six mood categories: anxiety-depression, fatigue, hostility, friendliness, energy, and happiness-satisfaction. Category scoring, however, was tedious, time-consuming, and not all the factors were sensitive to sleep loss.

We, therefore, constructed our own mood score, in such a way that it was maximally sensitive to sleep loss. The first step was to decide a priori which items were "positive" (e. g., satisfied, cheerful, efficient-all of which were predicted to decrease during sleep loss), and which items were "negative" (e. g., sleepy, unhappy, depressed--all of which were predicted to increase during sleep loss). Of the 52 items, 21 were judged positive and 31 were judged negative. We then tabulated the sum of the positive items and the sum of the negative items. A test for trend during the four baseline days showed no significant increase or decrease in the positive or negative scores. The baseline scores were therefore combined into an average positive score and an average negative score. The positive and negative sums were then tested for sensitivity to sleep loss by subtracting the baseline average score from the score after one night of sleep loss.

Of the 31 negative items, 21 increased as predicted, but the increase over baseline was not significant. Nineteen of the 21 positive items decreased after one night of sleep loss, and this decrease from the baseline average was significant ( $\bar{D}$  = -5.0; s = 7.4; N = 12). Additional analyses showed that P, the sum of the sensitive 19 positive items, was a better measure of one night of sleep loss than any weighted combination of the positive and negative items. These 19 positive items (active, alert, carefree, cheerful, able to concentrate, considerate, dependable, efficient, friendly, full of pep, good-natured, happy, kind, lively, pleasant, relaxed, satisfied, able to think clearly, able to work hard) were therefore used as the mood score in Experiment 2.

....

Tel Bila Latin

At the .05 level of significance, the statistical power of the Mood Scale is such that for 8 Ss, there is a 50% chance of finding a significant sleep loss effect; for 25 Ss, there is a 95% chance of finding a significant effect.

## Cross-Validation Study

The fourteen naval recruits in Experiment 2 (ages 18-21) were tested for four baseline days, and then underwent three nights of either stage 4 deprivation (N=7) or stage REM deprivation (N=7). The sleep stage deprivation was followed by one night of total sleep loss, and two recovery nights. (For a complete description of tests and methods used, see Johnson et al., 1974).

#### RESULTS

Student's <u>t</u>, backed by Wilcoxon's rank tests, was used for all tests of significance.

After one night of total sleep loss (plus 3 nights of differential sleep stage deprivation), both groups showed similar and significantly lower P scores, with an average drop of 7.3 points (s = 6.3; p < .001). Both groups returned to the baseline level after the first recovery sleep. The results are summarized in Table 1.

Table 1 about here) There were no significant differences between the REM-deprived and stage 4-deprived groups on any day. Both groups received lower P scores after three nights of sleep stage deprivation; the decrement from baseline was significant for the REM-deprived group. The sleep loss incurred during stage 4 deprivation was minimal. For the REM-deprived Ss, however, there was a cumulative sleep loss of about 200 min by the end of the third deprivation night. Thus, our Mood Scale was sensitive to a relatively small amount of sleep loss such as that accumulated during REM deprivation.

#### DISCUSSION

Subjective changes in mood after total sleep loss can be more sensitive than performance deficit or changes in quantitative indices of sleep behavior. Examination of the data of Experiment 1 (Lubin et al., 1974), showed that the Mood Scale, which requires about a minute to complete, was more sensitive to one night of sleep loss than the 20-min Word Memory Test or the 60-min X Crossout Test, but less sensitive than a 50-min counting task, a 60-min addition test, or a 50-min auditory vigilance test.

While the NPRU Mood Scale was designed to assess the effects of experimentally imposed sleep loss, clearly it can measure sleep loss regardless of cause and may offer an easily administered yet sensitive technique for assessing sleep loss effects in any context. Another important feature of the NPRU Mood Scale is that it lacks the learning factors common to many performance tests and therefore can be given repeatedly without showing practice effects.

In a study of the effects of one night of sleep loss on the structure of mood, Hendrick and Lilly (1970) administered a 44-item mood scale to college students and found that the items connoting negative feelings were as sensitive to sleep loss as the positive items. The negative items were not sensitive to sleep loss for our naval recruits. The 15 Army enlisted men in the Murray et al. study (1958) did not admit to feelings of fatigue or sleepiness after one night of sleep loss, even though experimental observers reported significant increases for these Ss on a sleepiness scale and a fatigue scale. These results indicate that Ss drawn from a military population are less likely than college students to admit increased feelings of fatigue, tension, or anger.

(Fig. 1 about here)

To broaden its applicability, the NPRU Mood Scale was buffered with ten negative items from Experiment 1 and from the Hendrick and Lilly study. These negative adjectives (annoyed, defiant, drowsy, dull, grouchy, jittery, sleepy, sluggish, tense, tired) were sensitive to sleep loss for one or both studies. Fig. 1 gives the final form of the NPRU Mood Scale.

Scoring. Each of the four possible response categories is assigned a weight: "not at all," 0; "a little," 1; "quite a bit," 2; "extremely," 3.

The sum of the responses to the 19 positive items is the P score. The sum of the responses to the 10 negative items is tabulated in the same way to obtain the N score. Combining the P and N scores is not recommended since the sensitivity of the N score to sleep loss is not conclusive.

#### **Footnotes**

In this paper, significance means .05 level or better, one-tail test, unless otherwise noted.

#### References

- Bohlin, G., & Kjellberg, A. Self-reported arousal during sleep deprivation and its relation to performance and physiological variables.

  Scandinavian Journal of Psychology, 1973, 14, 78-86.
- Hendrick, C., & Lilly, R. S. The structure of mod: A comparison between sleep deprivation and normal wakefulness conditions.

  Journal of Personality, 1970, 38, 453-465.
- Hoddes, E., Dement, W., & Zarcone, V. The history and use of the Stanford Sleepiness Scale. Psychophysiology, 1972, 9, 150. (Abstract)
- Hoddes, E., Zarcone, V., Smythe, H., Phillips, R., & Dement, W. C.

  Quantification of sleepiness: a new approach. Psychophysiology,
  1973, 10, 431-436.
- Johnson, E. III, & Myers, T. I. The development and use of the Primary

  Affect Scale (PAS). Naval Medical Research Institute, Bethesda, Md.,

  Research Report No. 31, 1967.
- Johnson, L. C., Naitoh, P., Moses, J. M., & Lubin, A. Interaction of REM deprivation and stage 4 deprivation with total sleep loss:

  Experiment 2. Psychophysiology, 1974, 11,...
- Lubin, A., Moses, J. M., Johnson, L. C., & Naitoh, P. The recuperative effects of REM sleep and stage 4 sleep on human performance after complete sleep loss: Experiment 1. Psychophysiology, 1974, 11,
- McNair, D. M., Lorr, M., & Droppleman, L. F. Profile of Mood States.

  Educational and Industrial Testing Service, San Diego, California,
  1971.
- Murray, E. J., Williams, H. L., & Lubin, A. Body temperature and psychological ratings during sleep deprivation. <u>Journal of Experimental Psychology</u>, 1958, <u>23</u>, 313-318.

- Spielberger, C. D. The Spielberger state-trait anxiety inventory.

  Palo Alto, Cal.: Consulting Psychologists Press, 1968.
- Thayer, R. E. Measurement of activation through self-report.

  Psychological Reports, 1967, 20, 663-678.
- Thayer, R. E. Activation states as assessed by verbal report and four psychophysiological variables. <a href="Psychophysiology">Psychophysiology</a>, 1970, 7, 86-94.
- Thayer, R. E. Studies of controlled self-reports of activation.

  Terminal Progress Report, National Institute of Mental Health,

  Public Health Service, MH-14248-01, 1971.

TABLE 1 P Scores a on the NPRU Mood Scale

Group	Baseline 1 2			Stage Dep.	Total Sleep Loss	Recover	y 2
REM- deprived (N=7)	M= SD=	34.6 8.4	32.3 6.1	29.4*	25.9* 9.5	32.0 9.7	35.1 10.1
S4- deprived (N=7)	M= SD=	31.9 4.6	29.9 7.7	27.9 6.2	23.9** 6.0	29.6 6.0	28.4 7.5

<sup>&</sup>lt;sup>a</sup>P scores range from 0 (extremely sleepy) to 57 (extremely active)  $\frac{a}{p}$  < .05

p < .01

NPRU MOCD SCALE 11ND-COM-6250 2 (5-73)

Instructions: For each item, choose one of the four answers that best describes how you feel <u>now</u>. Then put an "X" in that box.

NAME						AGE	SEX	DATE	
ITEM	NOT AT ALL	A LITTLE	QUITE A BIT	EXTREMELY	ITEM	NOT AT ALL	A LITTLE	QUITE A BIT	EXTREMELY
ACTIVE					GOOD-NATURED				
ALERT					GROUCHY				
ANNOYED					HAPPY				
CAREFREE					JITTERY				
CHEERFUL					KIND				
ABLE TO CONCENTRATE					LIVELY				
CONSIDERATE					PLEASANT				
DEFIANT					RELAXED				
DEPENDABLE					SATISFIED				
DROWSY					SLEEPY				
DULL					SLUGGISH				
EFFICIENT					TENSE				
FRIENDLY					ABLE TO THINK CLEARLY				
FULL OF PEP					TIRED				*
DO NOT FILL-IN SCOPES: N P				ABLE TO					

Fig. 1

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

S/N 0102 LF 014 6601

REPORT DOCUMENTATION PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM				
74-25 2. GOVT ACCESSION NO	3. RECIPIENT'S CATALOG NUMBER				
4 TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED				
Subjective Evaluation of the Effects of Sleep Loss: The NPRU Mood Scale	6. PERFORMING ORG. REPORT NUMBER				
Julie Moses, Ardie Lubin, Paul Naitoh	National Science Foundati Grants GB-6008 & BG-14829				
Naval Health Research Center San Diego, CA 92152  17	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS  MF15_525_004-9008DA5G				
Naval Medical Research & Development Command Bethesda, MD 20014	12. REPORT DATE 1974  13. NUMBER OF PAGES				
14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office)	15. SECURITY CLASS. (of this report)				
Bureau of Medicine & Surgery Department of the Navy	UNCLASSIFIED				
Washington, DC 20372	15a. DECLASSIFICATION DOWNGRADING SCHEDULE				
7. DISTRIBUTION STATEMENT (of the ebetrec! entered in Block 20, if different fro	m Report)				
8. SUPPLEMENTARY NOTES					
Adjective check list  Mood scale  Sleep loss  Task performance					
A mood scale consisting of 52 adjectives and ph naval recruits to determine which items were sensi of sleep loss. The items which were maximally sen then used to measure the effects of one night of t group of 14 young naval recruits. The two experim 29 adjectives and short phrases, the NPRU Mood Scaeasily, briefly, and repeatedly employed to detect to assist in evaluating task performance decrement	tive to at lteast one night sitive to sleep loss were otal sleep loss in another ents resulted in a list of le. This scale can be cumulative sleep loss and				
D I JAN 73 1473 EDITION OF I NOV 65 IS OBSOLETE	UNCLASSIFIED				

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)